

Acute HIV infection: the impact of anti-retroviral treatment on cellular immune responses

OTHER ARTICLE PUBLISHED IN THIS MINI-REVIEW SERIES ON INFECTIOUS DISEASE

Congenital cytomegalovirus infection: 50 years on. Clin Exp Immunol 2007; 149: doi:10.1111/j.1365-2249.2007.03454.x

C. B. Hicks,* C. Gay[†] and G. Ferrari[‡]

Duke University Medical Center, Department of

*Medicine and [†]Surgery, Durham, North

Carolina, USA, and [‡]University of North

Carolina at Chapel Hill, Center for Infectious

Diseases, Chapel Hill, NC, USA

Summary

The overall value of initiating anti-retroviral therapy during the acute phase of human immunodeficiency virus type 1 (HIV-1) infection remains unclear. From a clinical perspective, the lack of data from controlled randomized clinical trials limits understanding of long-term effects of treatment on the clinical course of HIV infection. Based on available data, the impact of anti-retroviral therapy during acute infection on the immune response against HIV-1 is not particularly encouraging. Recent observations on the very early depletion of lymphocyte reservoirs in the gastrointestinal tract may partially explain the limited benefit of anti-retroviral therapy initiated during the acute phase of HIV-1 infection. This may also help to explain the dichotomy between early observations demonstrating apparent immunological benefit with early anti-retroviral treatment that were associated none the less with inability to control viral replication following treatment interruption.

Keywords: acute infection, anti-retroviral therapy, cellular mediated immunity, HIV-1, immune reconstitution

Accepted for publication 17 May 2007

Correspondence: Guido Ferrari MD, Duke

University Medical Center, Department of

Surgery, PO Box 2926, Durham, North Carolina

27710, USA.

E-mail: gflmp@duke.edu

Introduction

Human immunodeficiency virus type 1 (HIV-1) can infect exposed individuals through mucosal surfaces or due to direct inoculation into the bloodstream. The virus subsequently infects CD4⁺ T lymphocytes directly or as a consequence of their interaction with virus-bearing dendritic cells in the lymphoid tissues [1,2]. As with many other viral infections, the onset of clinical manifestations in HIV-infected individuals is usually observed 2–4 weeks after exposure [3]. It has been reported that up to 90% of patients with acute HIV-1 infection (AHI) report to a physician due to the presence of symptoms that are most often described as a flu-like syndrome. The symptoms and signs seen in association with acute HIV infection may include fever, lymphadenopathy, pharyngitis, mucocutaneous lesions, myalgia/arthritis, headache, gastrointestinal symptoms (diarrhoea, nausea/vomiting) and weight loss [4]. Notably, none of these findings can be considered specific, although their persistence for periods as long as 10 weeks and the presence of mucocutaneous ulcers are suggestive of AHI [5]. A particular challenge in diagnosing AHI is the fact that the onset of symptoms

coincides typically with peak HIV viraemia [6], and usually precedes by 2–3 weeks the development of HIV-specific antibodies that are necessary for diagnosis using routine HIV screening tests [7]. Therefore, the diagnosis of AHI is often quite delayed (as long as 8–12 weeks after infection has occurred), and this delay probably has a great impact on the damage that occurs to the immune system before the initiation of anti-retroviral therapy (ART). In addition, delays in the diagnosis of AHI also prevent the implementation of secondary prevention measures with a group of patients who are at high risk of transmitting HIV during unprotected sexual encounters, due at least in part to the high levels of viraemia characteristic of this stage of infection [8,9].

Once the diagnosis has been made, additional challenges include the absence of clear guidelines on whether to initiate anti-retroviral treatment at this stage and the selection of an initial regimen if treatment is elected. Published guidelines for anti-retroviral therapy specifically note the lack of consensus in this area. The US Department of Health and Human Services (DHHS) guidelines (available at <http://aidsinfo.nih.gov/contentfiles/AdultandAdolescentGL.pdf>), developed by the working group of the office of AIDS

Research Advisory Council, and International AIDS Society (IAS)–USA guidelines [10] are appropriately vague, because of insufficient data to establish clear benefits of ART in acute infection [11]. The British HIV Association (BHIVA) recommends treatment for people with AHI for relief of symptoms only. As a result, the decision to initiate ART is left to the individual AHI patient and his or her health care provider. Common laboratory parameters used to guide treatment decisions for people with chronic HIV infection (such as HIV RNA levels and CD4 counts) are not clearly of value for acutely infected patients. As for choice of treatment regimen, most available data is on the use of standard-of-care regimens for chronic HIV infection, i.e. recommended first-line regimens for treatment-naïve patients as outlined in IAS–USA as well as BHIVA guidelines [10].

These various considerations illustrate the clinical challenges associated with diagnosing AHI and initiating ART treatment in a timely manner using an appropriate drug regimen. Bearing these factors in mind, we will discuss what is known of the impact current ART initiated during AHI may have on the reconstitution of cellular-mediated immune responses.

Impact of ART on cellular immune responses

The ability of HIV-1 to infect and to deplete the CD4⁺ T lymphocyte subset has been considered for years to be the hallmark of the pathogenesis of the acquired immunodeficiency syndrome (AIDS) [12]. The ongoing depletion of this pool of cells is thought to be the main cause of the progressive impairment of the adaptive and innate immune responses, the inability to control virus replication and the onset of opportunistic infections and other AIDS-related clinical manifestations.

Several studies have demonstrated that cellular-mediated immune responses may play a central role in controlling the initial burst of virus replication [13–15]. For years, CD8⁺ T cell responses have been considered the main line of defence during acute HIV infection and have been associated with delayed disease progression [16].

Early observations by Rosenberg and collaborators in a cohort of 10 untreated acutely infected individuals suggested that detectable CD4⁺ T cell responses directed mainly against the Gag p24 antigenic region were correlated inversely with subsequent virus load [17]. Interestingly, the study reported that three individuals who started ART at the time of AHI diagnosis and before seroconversion were also able to develop a strong anti-p24 lymphoproliferative response in synchrony with a decrease in viral load. This was in striking contrast with previous observations showing that chronically infected individuals lack strong lymphoproliferative responses to HIV-1 antigens, despite suppressive anti-retroviral therapy [18,19]. Subsequently, one study compared the magnitude and breadth of T cell responses in three groups of individuals: (1) those receiving ART

pre-seroconversion ($n = 19$); (2) those treated post-seroconversion and within 180 days of infection ($n = 11$); and (3) those treated during chronic infection more than 180 days after infection ($n = 10$) [20]. After receiving ART for 1 years, the pre-seroconversion group had the highest significant detectable CD4 lymphoproliferative responses [stimulation index (SI) = 43 ± 13] compared to either the post-seroconversion (SI = 29 ± 10) or the chronically infected groups (SI = 8 ± 3). Oxenius and collaborators observed similar beneficial effects from the initiation of ART during acute infection on the CD4⁺ T cell compartment. Their data indicated that the frequency of interferon (IFN)- γ -secreting CD4⁺ T cells was elevated in six patients up to 36 months post-treatment and directed mainly against HIV-1 Env and Gag antigens [21]. Moreover, Cossarizza and colleagues studied 18 individuals who received therapy during AHI for 1 year and compared the profile of their CD4⁺ V β T cell receptor repertoire to that observed in ART-treated chronically infected individuals and healthy non-infected donors [22]. The clonality of the CD4⁺ T cell population was apparently restored in AHI patients, but not in the chronically infected individuals. These findings, again, suggested a beneficial impact of ART initiated during AHI on the CD4⁺ T cell population.

Several studies have also reported that the initiation of ART during acute infection can boost CD8⁺ T cell responses [20,21,23–25]. These data suggested that CD8⁺ T cell responses were preserved by the initiation of therapy during AHI. However, the frequency and breadth of cytotoxic and IFN- γ -secreting CD8⁺ T cells decreased over time with suppression of viraemia, probably as a result of the lower antigen stimulation. Another encouraging observation was that the virus diversification in patients treated during AHI was lower and therefore escape mutants had not been generated, despite the strong cellular-mediated immune responses [20].

Although the initial observations on the importance of ART initiation in preserving T cell responses in acute infection was demonstrated in patients treated prior to seroconversion [20], more recent studies suggest that ART may have similar positive effects if initiated during the first 90 days post-infection [25,26]. Furthermore, recent investigations by Younes and colleagues on the breadth and magnitude of CD4⁺ T cell responses by lymphoproliferation and class-II-tetramer assays suggest that initiation of ART prior to seroconversion may impact negatively the maturation of HIV-specific IFN- γ -secreting CD4⁺ T cells as a consequence of the limited time for interaction of antigen-presenting cells with virus due to low antigenaemia [26].

With respect to duration of therapy, Oxenius and collaborators suggested that even a short course of ART treatment initiated during AHI and continued for up to 24 weeks post-infection could exert beneficial effects on the longevity and breadth of CD8⁺ and CD4⁺ T cell responses [21].

Impact of T cell responses generated under AHI-ART on virus load

A consistent finding among individuals treated during AHI has been higher proliferative capacity and frequency and breadth of IFN- γ - and interleukin (IL)-2-secreting T cells in the groups receiving ART during AHI. This outcome was observed independently of the duration of therapy, ranging from 12 weeks [27] to 98 weeks [28]. Based on these findings and those outlined in the previous section, indicating that ART initiation within 90 days of infection preserved strong T cell responses, one can hypothesize that subsequent treatment interruption in these subjects might result in control of HIV infection. Although viral control was demonstrated in three of 14 individuals studied by Rosenberg and colleagues [17], subsequent studies did not observe consistent viral control following treatment interruption. Therefore, despite preservation of cellular immune responses, initiation of ART during AHI has not shown a consistent and significant impact on decreasing viral rebound following treatment interruption [27–30].

In summary, there are no clear data indicating that early initiation of ART during AHI lowers the viral set-point or delays disease progression following treatment interruption.

Controlling virus replication: a multi-faceted picture

The simplest potential reason for the inability to demonstrate control of virus replication following ART interruption in patients treated during AHI is that the focus on cellular-mediated immune responses as the key factor in controlling HIV-1 replication may be incorrect. Indeed, this relationship may not be central, and other components of the immune response against HIV-1, as well as genetic factors, could correlate with protection from infection and/or disease progression [31].

If an overly narrow focus on cellular responses is a limiting factor in understanding viral control, consideration of both humoral and innate immunity may be important. Both responses can be influenced positively by ART initiated during AHI [32,33], but these factors may still not be sufficient for control of virus replication following treatment interruption.

As for other factors, even large cohort studies have been unable to establish whether genetic factors, such as selected HLA alleles and killer inhibitory receptor (KIR) haplotypes, may contribute to the control of virus replication [34–36] or whether they may have synergistic effects with the initiation of ART during AHI.

ART-preserved cellular immunity: reasons for failure

Lack of a clear understanding regarding which factors are important in control of viral replication limits the ability to

discern which parameters should be studied to assess the effect of ART initiated during AHI [31]. Consideration of data on T cell responses in individuals who control HIV-1 replication successfully without ART for up to 18 years post-infection is illuminating. Viral control in these 'long-term non-progressor' patients (LTNP) may not relate to a single factor such as proliferative capability and/or single cytokine production. Migueles and colleagues demonstrated that the preservation of proliferative capability and perforin production of CD8⁺ T lymphocytes set LTNP apart from those who do not control virus replication [37]. Betts and collaborators recently reported a higher frequency and breadth of CD8⁺ T cell responses with a multi-functional profile in LTNP compared with subjects who do not control virus replication successfully [38]. Both studies revealed that the quality of cellular-mediated immune responses may be more important than their mere frequency or breadth. One could argue that even these parameters simply reflect the preservation of CD4⁺ T cell responses that sustain other cellular, humoral and innate components of the anti-viral responses. However, the question of how cellular immune responses are preserved in a limited number of infected subjects, and not in others, remains.

One hypothesis is that the immune responses involved in control of virus replication after treatment interruption may be compared to those described as a recall response after antigen-priming of the immune system, although virus replication and therefore antigen stimulation still occur during the latent phase of HIV-1 infection [12,39]. Janssen *et al.* demonstrated the importance of CD4⁺ T cells providing the correct signal to the CD8⁺ T cell compartment during the priming of the immune responses, as the recall CD8⁺ T responses are impaired in the absence of this interaction [40]. In the course of HIV infection, the maturation and the functionality of CD8⁺ T cells are impaired [41–45], reminders that the two fundamental effects of HIV infection on the immune system are severe depletion of CD4⁺ T lymphocytes and persistent cellular activation. Accordingly, this raises the question: 'is ART initiated during AHI effective in preventing loss of CD4⁺ T lymphocytes that may be one of the key factors in correctly priming the immune system?'

Loss of CD4⁺ T cells during AHI

The preferential infection of anti-HIV-specific memory CD4⁺ T cells during the course of HIV-1 infection has been well documented [46,47]. Surprisingly, it has also been shown that the CCR5⁺CD4⁺ lymphocytes that reside in the lamina propria of the lower gastrointestinal tract are the major targets for both simian immunodeficiency virus (SIV) and HIV-1 replication, independently of the route of infection [48–53]. These studies demonstrated that up to 60% of CCR5⁺CD4⁺ cells can be infected at the peak of

viraemia, and 80% are depleted within 4 weeks from infection as a consequence of direct virus replication [51] or apoptosis [52,53]. It is probably this early and massive depletion of CD4⁺ T cell subsets that limits priming of the immune system and therefore control of HIV-1 infection during the early and late stages through functional cellular immune responses.

Recently, several groups have reported on the effect of ART initiated during AHI on reconstitution of mucosal lymphoid tissue. Results from these studies have been consistently disappointing and revealed a lack of reconstitution of lamina propria lymphocytes. In fact, Mehandru and collaborators studied 18 subjects, nine of whom started therapy pre-seroconversion and continued treatment for up to 7 years post-diagnosis [54]. Serial gastrointestinal biopsies were performed in a subset of these subjects at 1, 2 and 3 years' post-treatment initiation. Although reconstitution of CD4⁺ T cells in the peripheral blood reached 67% at 3 years, the percentage of CD4⁺ T cells in the mucosa of the gastrointestinal tract never exceeded 45% of the level observed in healthy individuals. Moreover, immunohistochemistry analysis of the biopsies revealed that the immune-inductive sites were fully reconstituted, whereas the effector sites of the lamina propria were persistently depleted. This partial reconstitution of the CD4⁺ and CD8⁺ T cell population in the mucosal tract confirmed what had previously been observed by Guadalupe and collaborators in a smaller number of individuals ($n = 3$) during the first 14 months of therapy initiated following seroconversion [55]. An additional interesting finding by Mehandru and his group was that activated CD4⁺ and CD8⁺ T cells were still elevated at the mucosal sites after 3–7 years of treatment, even in the presence of a minimal number of cells expressing viral mRNA. It has been postulated that the incomplete reconstitution of resident T cell populations in the lamina propria may allow a continuous leakage of foreign antigens from the lumen of the gastrointestinal tract that can persistently activate T cells and predispose them to apoptosis [54,56]. One premise behind the use of ART is that a high level of virus replication is the main reason for CD4⁺ T cell depletion. However, it has been suggested recently that this may not be the case and that other co-factors may play an important role. For example, persistent antigenic stimulation may be a critical factor in the pathogenesis of damage to mucosal surfaces which, in turn, leads to apoptosis of lymphocytes [57]. In this situation, these secondary effects on the immune system cannot be addressed by ART therapy alone.

Taken together, these studies indicate that massive depletion of the T lymphocyte pool at the mucosal level takes place before it is possible to diagnose AHI in most patients, and certainly prior to initiation of ART. This lag probably impedes reconstitution or recovery of mucosal T cells, and could be the major factor preventing recovery from immunological damage sustained during peak HIV-1 replication.

The inevitable consequence is, in most cases, that the immune system will not be able to control virus replication upon treatment interruption.

Clinical impact of ART

From a clinical perspective, the effect of ART on patient outcome may be more favourable overall than that observed in immunological studies. Hecht and collaborators detected a trend towards higher CD4 cell counts and lower HIV RNA levels among 13 people treated with ART during acute HIV infection when measured 24 weeks after ART interruption, if therapy was initiated within 2 weeks of seroconversion [27]. Initiation of therapy at a later time-point showed less benefit in 45 individuals when measured 72 weeks after treatment interruption. Moreover, there was a substantial decrease in virus load (approximating a two-thirds-log₁₀ reduction) observed in the adjusted analysis at 72 weeks among those who were treated earlier, a reduction which corresponded to an approximate 50% reduction in the risk of HIV disease progression over 6 years [58].

Despite the fact that ART initiated during AHI may not consistently alter the viral set-point following treatment interruption, we need to consider other therapeutic benefits which may still justify its use: (1) control of symptoms related to acute retroviral syndrome; (2) reduction in viral load which probably decreases the risk of transmission; (3) potential slowing of the rate of disease progression among patients with a genetic predisposition; (4) decrease in the size of the latent HIV pool; (5) possible control of viraemia and disease progression during which time advances in therapeutic vaccines and other treatment modalities may evolve; and (6) possible slowing of the pace of CD4 cell count decline prior to institution of long-term ART.

Therefore, despite the fact that initiation of ART during AHI may not have a significant impact on immune reconstitution as currently understood, it may alter clinical outcomes for those diagnosed during acute HIV-1 infection. As noted by others, findings which could determine the most predictive immunological parameters for favourable clinical outcome have yet to be demonstrated in a randomized clinical trial (RCT) [59]. A randomized controlled trial of treatment *versus* no treatment in the setting of recent HIV infection is currently ongoing (ACTG 5217), but will not be able to address the more probable benefit of ART initiated during 'acute' infection, as most enrolled patients will be outside the acute infection time-frame by the time therapy is initiated. Given the challenges of conducting a large randomized study in patients with AHI, it is not likely that convincing data will be available for guideline development in the foreseeable future. Thus, the clinical use of ART during AHI will continue to be based on the best available evidence and expert opinion. As such, it should still be carefully evaluated on a case-by-case basis.

Acknowledgements

This work was supported by grants from the National Institute of Health/NIAID: (RO1-AI-050483-06) (G. F.); 2K24 AI01608-08, U01 AI041530-08 (C. H.); U01 AI067854 (C. H. and CG).

References

- Wu L, Kewa VN, Ramani I. Dendritic-cell interactions with HIV: infection and viral dissemination. *Nat Rev Immunol* 2006; **6**:859–68.
- Morrow G, Vachot L, Vagenas P, Robbani M. Current concepts of HIV transmission. *Curr HIV/AIDS Rep* 2007; **4**:29–35.
- Ridzon R, Gallagher K, Ciesielski C *et al.* Simultaneous transmission of human immunodeficiency virus and hepatitis C virus from a needle-stick injury. *N Engl J Med* 1997; **336**:919–22.
- Niu MT, Stein DS, Schnittman SM. Primary human immunodeficiency virus type 1 infection: review of pathogenesis and early treatment intervention in humans and animal retrovirus infections. *J Infect Dis* 1993; **168**:1490–501.
- Kahn JO, Walker BD. Acute human immunodeficiency virus type 1 infection. *N Engl J Med* 1998; **339**:33–9.
- Fiebig EW, Wright DJ, Rawal BD *et al.* Dynamics of HIV viremia and antibody seroconversion in plasma donors: implications for diagnosis and staging of primary HIV infection. *AIDS* 2003; **17**:1871–9.
- Lindback S, Thorstensson R, Karlsson AC *et al.* Diagnosis of primary HIV-1 infection and duration of follow-up after HIV exposure. Karolinska Institute Primary HIV Infection Study Group. *AIDS* 2000; **14**:2333–9.
- Cohen MS, Pilcher CD. Amplified HIV transmission and new approaches to HIV prevention. *J Infect Dis* 2005; **191**:1391–3.
- Brenner BG, Roger M, Routy JP *et al.* High rates of forward transmission events after acute/early HIV-1 infection. *J Infect Dis* 2007; **195**:951–9.
- Hammer SM, Saag MS, Schechter M *et al.* Treatment for adult HIV infection: 2006 recommendations of the International AIDS Society–USA panel. *JAMA* 2006; **296**:827–43.
- Babiker A, Darbyshire J, Pezzotti P *et al.* Short-term CD4 cell response after highly active anti-retroviral therapy initiated at different times from seroconversion in 1500 seroconverters. *J Acquir Immune Defic Syndr* 2003; **32**:303–10.
- Pantaleo G, Graziosi C, Demarest JF *et al.* HIV infection is active and progressive in lymphoid tissue during the clinically latent stage of disease. *Nature* 1993; **362**:355–8.
- Borrow P, Lewicki H, Hahn BH, Shaw GM, Oldstone MBA. Virus-specific CD8+ cytotoxic T lymphocyte activity associated with control viremia in primary human immunodeficiency virus type 1 infection. *J Virol* 1994; **68**:6103–10.
- Koup RA, Safrit JF, Cao Y *et al.* Temporal association of cellular immune response with the initial control of viremia in primary human immunodeficiency virus type 1 syndrome. *J Virol* 1994; **68**:4650–5.
- Safrit JT, Andrews CA, Zhu T, Ho DD, Koup RA. Characterization of human immunodeficiency virus type 1-specific cytotoxic T lymphocyte clones isolated during acute seroconversion: recognition of autologous virus sequences within a conserved immunodominant epitope. *J Exp Med* 1994; **179**:463–72.
- Harrer T, Harrer E, Kalams SA *et al.* Cytotoxic T lymphocytes in asymptomatic long-term nonprogressing HIV-1 infection. *J Immunol* 1996; **156**:2616–23.
- Rosenberg ES, Altfeld M, Poon SH *et al.* Immune control of HIV-1 after early treatment of acute infection. *Nature* 2000; **407**:523–6.
- Plana M, Garcia F, Gallart T, Miro JM, Gatell JM. Lack of T-cell proliferative response to HIV-1 antigens after 1 year of highly active anti-retroviral treatment in early HIV-1 disease. Immunology Study Group Spanish EARTH-1 Study. *Lancet* 1998; **352**:1194–5.
- Rinaldo CR Jr, Liebmman JM, Huang XL *et al.* Prolonged suppression of human immunodeficiency virus type 1 (HIV-1) viremia in persons with advanced disease results in enhancement of CD4 T cell reactivity to microbial antigens but not to HIV-1 antigens. *J Infect Dis* 1999; **179**:329–36.
- Altfeld M, Rosenberg ES, Shankarappa R *et al.* Cellular immune responses and viral diversity in individuals treated during acute and early hiv-1 infection. *J Exp Med* 2001; **193**:169–80.
- Oxenius A, Price DA, Easterbrook PJ *et al.* Early highly active anti-retroviral therapy for acute HIV-1 infection preserves immune function of CD8+ and CD4+ T lymphocytes. *Proc Natl Acad Sci USA* 2000; **97**:3382–7.
- Cossarizza A, Poccia F, Agrati C *et al.* Highly active anti-retroviral therapy restores CD4+ Vbeta T-cell repertoire in patients with primary acute HIV infection but not in treatment-naive HIV+ patients with severe chronic infection. *J Acquir Immune Defic Syndr* 2004; **35**:213–22.
- Dalod M, Harzic M, Pellegrin I *et al.* Evolution of cytotoxic T lymphocyte responses to human immunodeficiency virus type 1 in patients with symptomatic primary infection receiving anti-retroviral triple therapy. *J Infect Dis* 1998; **178**:61–9.
- Alter G, Hatzakis G, Tsoukas CM *et al.* Longitudinal assessment of changes in HIV-specific effector activity in HIV-infected patients starting highly active anti-retroviral therapy in primary infection. *J Immunol* 2003; **171**:477–88.
- Lacabaratz-Porret C, Urrutia A, Doisme JM *et al.* Impact of anti-retroviral therapy and changes in virus load on human immunodeficiency virus (HIV)-specific T cell responses in primary HIV infection. *J Infect Dis* 2003; **187**:748–57.
- Younes SA, Trautmann L, Yassine-Diab B *et al.* The duration of exposure to HIV modulates the breadth and the magnitude of HIV-specific memory CD4+ T cells. *J Immunol* 2007; **178**:788–97.
- Hecht FM, Wang L, Collier A *et al.* A multicenter observational study of the potential benefits of initiating combination anti-retroviral therapy during acute HIV infection. *J Infect Dis* 2006; **194**:725–33.
- Jansen CA, De Cuyper IM, Steingrover R *et al.* Analysis of the effect of highly active anti-retroviral therapy during acute HIV-1 infection on HIV-specific CD4 T cell functions. *AIDS* 2005; **19**:1145–54.
- Kaufmann DE, Lichterfeld M, Altfeld M *et al.* Limited durability of viral control following treated acute HIV infection. *Plos Med* 2004; **1**:e36.
- Streeck H, Jessen H, Alter G *et al.* Immunological and virological impact of highly active anti-retroviral therapy initiated during acute HIV-1 infection. *J Infect Dis* 2006; **194**:734–9.
- Heeney JL, Plotkin SA. Immunological correlates of protection from HIV infection and disease. *Nat Immunol* 2006; **7**:1281–4.
- Robinson JE, Elliott DH, Martin EA, Micken K, Rosenberg ES. High frequencies of antibody responses to CD4 induced epitopes in HIV infected patients started on HAART during acute infection. *Hum Antibodies* 2005; **14**:115–21.

- 33 Alter G, Teigen N, Ahern R *et al.* Evolution of innate and adaptive effector cell functions during acute HIV-1 infection. *J Infect Dis* 2007; **195**:1452–60.
- 34 Gao X, Bashirova A, Iversen AK *et al.* AIDS restriction HLA allotypes target distinct intervals of HIV-1 pathogenesis. *Nat Med* 2005; **11**:1290–2.
- 35 Khakoo SI, Thio CL, Martin MP *et al.* HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science* 2004; **305**:872–4.
- 36 O'Brien SJ, Gao X, Carrington M. HLA and AIDS: a cautionary tale. *Trends Mol Med* 2001; **7**:379–81.
- 37 Migueles SA, Laborico AC, Shupert WL *et al.* HIV-specific CD8+ T cell proliferation is coupled to perforin expression and is maintained in nonprogressors. *Nat Immunol* 2002; **3**:1061–8.
- 38 Betts M, Nason MC, West SM *et al.* HIV nonprogressors preferentially maintain highly functional HIV-specific CD8+ T-cells. *Blood* 2006; **107**:4781–9.
- 39 Embretson J, Zupancic M, Ribas JL *et al.* Massive covert infection of helper T lymphocytes and macrophages by HIV during the incubation period of AIDS. *Nature* 1993; **362**:359–63.
- 40 Janssen EM, Lemmens EE, Wolfe T, Christen U, von Herrath MG, Schoenberger SP. CD4+ T cells are required for secondary expansion and memory in CD8+ T lymphocytes. *Nature* 2003; **421**:852–6.
- 41 Appay V, Nixon DF, Donahoe SM *et al.* HIV-specific CD8(+) T cells produce antiviral cytokines but are impaired in cytolytic function. *J Exp Med* 2000; **192**:63–76.
- 42 Champagne P, Ogg GS, King AS *et al.* Skewed maturation of memory HIV-specific CD8 T lymphocytes. *Nature* 2001; **410**:106–11.
- 43 Day CL, Kaufmann DE, Kiepiela P *et al.* PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature* 2006; **443**:350–4.
- 44 Petrovas C, Casazza JP, Brenchley JM *et al.* PD-1 is a regulator of virus-specific CD8+ T cell survival in HIV infection. *J Exp Med* 2006; **203**:2281–92.
- 45 Trautmann L, Janbazian L, Chomont N *et al.* Upregulation of PD-1 expression on HIV-specific CD8+ T cells leads to reversible immune dysfunction. *Nat Med* 2006; **12**:1198–202.
- 46 Douek DC, Brenchley JM, Betts MR *et al.* HIV preferentially infects HIV-specific CD4+ T cells. *Nature* 2002; **417**:95–8.
- 47 Douek DC, Picker LJ, Koup RA. T cell dynamics in HIV-1 infection. *Annu Rev Immunol* 2003; **21**:265–304.
- 48 Veazey RS, DeMaria M, Chalifoux LV *et al.* Gastrointestinal tract as a major site of CD4+ T cell depletion and viral replication in SIV infection. *Science* 1998; **280**:427–31.
- 49 Veazey RS, Gauduin MC, Mansfield KG *et al.* Emergence and kinetics of simian immunodeficiency virus-specific CD8(+) T cells in the intestines of macaques during primary infection. *J Virol* 2001; **75**:10515–9.
- 50 Brenchley JM, Schacker TW, Ruff LE *et al.* CD4+ T cell depletion during all stages of HIV disease occurs predominantly in the gastrointestinal tract. *J Exp Med* 2004; **200**:749–59.
- 51 Mehandru S, Poles MA, Tenner-Racz K *et al.* Primary HIV-1 infection is associated with preferential depletion of CD4+ T lymphocytes from effector sites in the gastrointestinal tract. *J Exp Med* 2004; **200**:761–70.
- 52 Li Q, Duan L, Estes JD *et al.* Peak SIV replication in resting memory CD4+ T cells depletes gut lamina propria CD4+ T cells. *Nature* 2005; **434**:1148–52.
- 53 Mattapallil JJ, Douek DC, Hill B, Nishimura Y, Martin M, Roederer M. Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV infection. *Nature* 2005; **434**:1093–7.
- 54 Mehandru S, Poles MA, Tenner-Racz K *et al.* Lack of mucosal immune reconstitution during prolonged treatment of acute and early HIV-1 infection. *Plos Med* 2006; **3**:e484.
- 55 Guadalupe M, Reay E, Sankaran S *et al.* Severe CD4+ T-cell depletion in gut lymphoid tissue during primary human immunodeficiency virus type 1 infection and substantial delay in restoration following highly active anti-retroviral therapy. *J Virol* 2003; **77**:11708–17.
- 56 Brenchley JM, Price DA, Schacker TW *et al.* Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med* 2006; **12**:1365–71.
- 57 Brenchley JM, Price DA, Douek DC. HIV disease: fallout from a mucosal catastrophe? *Nat Immunol* 2006; **7**:235–9.
- 58 Mellors JW, Munoz A, Giorgi JV *et al.* Plasma viral load and CD4+ lymphocytes as prognostic markers of HIV-1 infection. *Ann Intern Med* 1997; **126**:946–54.
- 59 Kinloch-de Loes S. Treatment of acute HIV-1 infection: is it coming of age? *J Infect Dis* 2006; **194**:721–4.